

The distribution of CO₂ between atmosphere, hydrosphere, and lithosphere; minimal influence from anthropogenic CO₂ on the global "Greenhouse Effect".

Tom V. Segalstad

Mineralogical-Geological Museum
University of Oslo
Sars' Gate 1, N-0562 Oslo
Norway

1. Introduction

It has recently been created a belief among people that an apparent increase in atmospheric CO₂ concentration is caused by anthropogenic burning of fossil carbon in petroleum, coal, and natural gas. The extra atmospheric CO₂ has been claimed to cause global climatic change with a significant atmospheric temperature rise, of 1.5 to 4.5°C in the next decennium (Houghton et al., 1990). This postulate is here discussed and rejected on energetic and geochemical grounds.

2. Heat energy and temperatures

Our relatively high global atmospheric temperature near the surface of the Earth, with an average of 14 to 15°C, is caused by heat-absorbing gases in the atmosphere, mainly H₂O vapor. Without the Earth's atmosphere the surface temperature would be approximately -18°C.

The Earth receives about 1368 W/m² of radiative heat from the Sun. The total amount of this heat withheld, approximately 11%, in the Earth's lower atmosphere, has traditionally been named the Earth's "Greenhouse Effect". For a cloudless atmosphere this effect is on the average about 146 W/m² for the Earth, with an uncertainty of ± 5 to 10 W/m² due to analytic uncertainties and natural climatic variations. All human activities have been claimed to contribute about 1.3% of this (approx. 2 W/m²), while a hypothetic doubling of the atmospheric CO₂ concentration would contribute about 2.6% (approx. 4 W/m²) to the present "Greenhouse Effect" (Raval & Ramanathan, 1989; Ramanathan et al., 1989).

150 year long time series of temperature measurements are covering too short time spans to be useful for climate prediction, in order to be used as "evidence" for anthropogenic heating (or cooling). The global mean temperature has risen and fallen several times over the last 400 years, with no evidence of anthropogenic causes, although strong explosive volcanic eruptions have caused periodically colder climates (Jaworowski et al., 1992 a).

It should also be noted that clouds can reflect up to approx. 50 W/m² and can absorb up to approx. 30 W/m² of the solar radiation (Ramanathan et al., 1989), making the Earth's average "Greenhouse Effect" vary naturally within approx. 96 and 176

W/m^2 . Hence the anticipated anthropogenic atmospheric CO_2 heat absorption is much smaller than the natural variation of the Earth's "Greenhouse Effect" (Segalstad & Jaworowski, 1991).

The oceans act as a huge heat energy buffer; the global climate is primarily governed by the enormous amount of heat stored in the oceans (total mass approx. 1.4×10^{24} g), rather than the minute amount of heat withheld in the heat-absorbing part of the atmosphere (total mass approx. 1.4×10^{18} g), a mass difference of one million times (Peixoto & Oort, 1992). Most of the atmospheric heat absorption occurs in water vapor (total mass approx. 1.3×10^{19} g), which is equivalent to a uniform layer of only 2.5 cm of liquid water covering the globe, with a residence time of about 9 days (Peixoto & Oort, 1992).

The total internal energy of the whole ocean is more than 1.6×10^{27} Joule, about 2000 times larger than the total internal energy 9.4×10^{23} Joule of the whole atmosphere. Note that this energy is defined with respect to 0 Kelvin (Peixoto & Oort, 1992).

Furthermore the cryosphere (ice sheets, sea ice, permafrost, and glaciers; total mass of the continental ice is approx. 3.3×10^{22} g) plays a central role in the Earth's climate as an effective heat sink for the atmosphere and oceans, with a large latent heat of melting on the order of 9.3×10^{24} Joule, a hypothetical energy equivalent to cooling the entire oceans by about $2^\circ C$ (5.8×10^{24} J/ $^\circ C$). For comparison, the energy needed to warm the entire atmosphere by $1^\circ C$ is only 5.1×10^{21} Joule (Oerlemans & van der Veen, 1984).

Hence it will be impossible to melt the Earth's ice caps and thereby increase the sea level just by increasing the heat energy of the atmosphere through a few percent by added heat absorption of anthropogenic CO_2 in the lower atmosphere.

3. CO_2 measurements in atmosphere and ice cores

Houghton et al. (1990) claim in their section 1.2.5 three evidences that the contemporary atmospheric CO_2 increase is anthropogenic: *First*, CO_2 measurements from ice cores show a 21% rise from 280 to 353 ppmv (parts per million by volume) since pre-industrial times; *second*, the atmospheric CO_2 increase closely parallels (sic!) the accumulated emission trends from fossil fuel combustion and from land use changes, although the annual increase has been smaller each year than the fossil CO_2 input [some 50% deviation, e.g. Kerr, 1992]; *third*, the isotopic trends of ^{13}C and ^{14}C agree qualitatively (sic!) with those expected due to the CO_2 emissions from fossil fuels and the biosphere.

Jaworowski et al. (1992 a, 1992 b) reviewed published CO_2 measurements from ice cores, and emphasized that the pre-industrial atmospheric CO_2 concentration, according to early accurate analyses, was many times larger (measurements up to 2450 ppmv) than the present atmospheric value. They also pointed out that CO_2 variations in ice is mainly an artificial effect of a large number of natural physical-chemical processes in ice and the recovered ice cores. These effects dominate over the eventual traces of anthropogenic CO_2 . Criticism of the methodology has also independently been presented by Heyke (1992 a, 1992 b, 1992 c).

Jaworowski et al. (1992 a) have presented a number of criticisms regarding the methodology of atmospheric CO₂ measurements, including spectroscopic instrumental peak overlap errors (from N₂O, CH₄, and CFCs in the air). They also pointed out that the CO₂ measurements at current CO₂ observatories use a procedure involving a subjective editing (Keeling et al., 1976) of measured data, only representative of a few tens of percent of the total data. There are also fundamental problems connected with the use of stable carbon isotopes (¹³C/¹²C) in tree rings for model calculations of earlier atmospheres' CO₂ concentration, a method which now seems to have been abandoned (Jaworowski et al., 1992 a).

The third evidence, based on carbon isotopes, will be discussed below in Section 5.

4. Chemical laws for distribution of CO₂ in nature


Statistically it has been found that the atmospheric CO₂ concentration rises after temperature rises (Kuo et al., 1990), and it has been suggested that the reason is that cold water dissolves more CO₂ (e.g. Segalstad, 1990). Hence, if the water temperature increases, the water cannot keep as much CO₂ in solution, resulting in CO₂ degassing from the water to the atmosphere. According to Takahashi (1961) heating of sea water by 1°C will increase the partial pressure of atmospheric CO₂ by 12.5 ppmv during upwelling of deep water. For example 12°C warming of the Benguela Current should increase the atmospheric CO₂ concentration by 150 ppmv.

Volk & Liu (1988) modelled the CO₂ flux between atmosphere and oceans, and concluded that approximately 70% of the flux was governed by this "thermal solubility pump", while approximately 30% was governed by the organic nutrient "biological pump". Faure (1990) estimated that ca. 4000 GT (Gigatonnes = billion metric tonnes) of CO₂ is transferred by degassing of the ocean via the atmosphere to the continental biosphere from the end of a glaciation to an interglacial stage.

From a geochemical consideration of sedimentary rocks deposited throughout the Earth's history, and the chemical composition of the ocean and atmosphere, Holland (1984) showed that degassing from the Earth's interior has given us chloride in the ocean; and nitrogen, CO₂, and noble gases in the atmosphere. Mineral equilibria have established concentrations of major cations and H⁺ in the ocean, and the CO₂ concentration in the atmosphere, through different chemical buffer reactions. Biological reactions have given us sulphate in the ocean and oxygen in the atmosphere.

Carbon dioxide is an equally important requisite for life on Earth as oxygen. Plants need CO₂ for their living (the photo synthesis), and humans and animals breath out CO₂ from their respiration. In addition to this biogeochemical balance, there is also an important geochemical balance. CO₂ in the atmosphere is in equilibrium with carbonic acid dissolved in the ocean, which in term is close to CaCO₃ saturation and in equilibrium with carbonate shells of organisms and lime (calcium carbonate; limestone) in the ocean through the following reactions (where *s* indicates the solid state, *aq* is the aqueous state, and *g* is the gaseous state):

Partial reactions:



Net reaction:



In addition there are a number of different aqueous metal complexes of lesser concentrations.

A buffer can be defined as a reaction system which modifies or controls the value of an intensive (i.e. mass independent) thermodynamic variable (pressure, temperature, concentration, pH, etc.). Our carbonate system above will act as a pH buffer, by the presence of a weak acid (H_2CO_3) and a salt of the acid (CaCO_3). The concentration of $\text{CO}_2 (g)$ and of $\text{Ca}^{2+} (aq)$ will in the equilibrium Earth system also be buffered by the presence of CaCO_3 , at a given temperature. **If the partial pressure of $\text{CO}_2 (g)$ is increased, the net reaction will go towards the right because of the Law of Mass Action.** If the temperature changes, the chemical equilibrium constant will change, and move the equilibrium to the left or right. The result is that the partial pressure of $\text{CO}_2 (g)$ will increase or decrease. The equilibrium will mainly be governed by Henry's Law: the partial pressure of $\text{CO}_2 (g)$ in the air will be proportional to the concentration of $\text{CO}_2 (aq)$ dissolved in water. **The proportional constant is the Henry's Law Constant,** which is strongly temperature dependent, and lesser dependent on total pressure and salinity (Drummond, 1981).

Questions have been raised about how strong this buffer is. It has been postulated (Bolin & Keeling, 1963) that an increase in atmospheric CO_2 will be balanced when only approximately one tenth of this is dissolved in the ocean. This postulate fails for a number of reasons. An increase in atmospheric CO_2 will namely increase the buffer capacity of ocean water, and thereby strengthen the ocean's capacity to moderate an increase of atmospheric CO_2 ; maximum buffer capacity for the system $\text{CO}_2 - \text{H}_2\text{O}$ is reached at 2.5 to 6 times the present atmospheric partial pressure of CO_2 , depending on temperature and alkalinity (Butler, 1982). According to Maier-Reimer & Hasselmann (1987) the borate system also increases the ocean storage capacity for CO_2 by more than 20% over an ocean with the carbonate-system alone.

Furthermore, this carbonate buffer is not the only buffer active in the atmosphere / hydrosphere / lithosphere system. **The Earth has a set of other buffering mineral reactions.** The geochemical equilibrium system anorthite $\text{CaAl}_2\text{Si}_2\text{O}_8$ - kaolinite $\text{Al}_2\text{Si}_2\text{O}_5(\text{OH})_4$ has by the pH of ocean water a buffer capacity which is thousand times larger than a 0.001 M carbonate solution (Stumm & Morgan, 1970). In addition we have clay mineral buffers, and a calcium silicate + $\text{CO}_2 \leftrightarrow$ calcium carbonate + SiO_2 buffer (MacIntyre, 1970; Krauskopf, 1979). These buffers all act as a "security net" under the most important buffer: $\text{CO}_2 (g) \leftrightarrow \text{HCO}_3^- (aq) \leftrightarrow \text{CaCO}_3 (s)$. All together these buffers give in principle an **infinite buffer capacity** (Stumm & Morgan, 1970).

Stable carbon isotopes ($^{13}\text{C}/^{12}\text{C}$) show that CO_2 in the atmosphere is in chemical equilibrium with ocean bicarbonate and lithospheric carbonate (Ohmoto, 1986). The chemical equilibrium constants for the chemical reactions above provide us with a partition coefficient for CO_2 between the atmosphere and the ocean of approximately 1 : 50 (approx. 0.02) at the global mean temperature (Revelle & Suess, 1957; Skirrow, 1975). This means that for an atmospheric doubling of CO_2 , there will have to be supplied 50 times more CO_2 to the ocean to obtain chemical equilibrium. This total of 51 times the present amount of atmospheric CO_2 carbon is more than the known reserves of fossil carbon. It is possible to exploit approximately 7000 GT of fossil carbon, which means, if all this carbon is supposed to be burned, that the atmospheric CO_2 can be increased by 20% at the most under geochemical equilibrium at constant present surface temperature.

^{14}C isotopes show that the circulation time for carbon in the upper part of the ocean is some few decades (Druffel & Williams, 1990). This is sufficient time for the ocean to absorb an increase in atmospheric CO_2 from burning of fossil fuel at the present projected rate (Jaworowski et al., 1992 a).

5. Carbon isotopes in atmospheric CO_2

Houghton et al. (1990) assumed for the IPCC model 21% of our present-day atmospheric CO_2 has been contributed from burning of fossil fuel. This has been made possible by CO_2 having a "rough indication" (sic!) lifetime of 50 - 200 years. It is possible to test this assumption by inspecting the stable $^{13}\text{C}/^{12}\text{C}$ isotope ratio (expressed as $\delta^{13}\text{C}_{\text{PDB}}^*$) of atmospheric CO_2 . It is important to note that this value is the net value of mixing all different CO_2 components, and would show the results of all natural and non-natural (i.e. anthropogenic) processes involving CO_2 .

The natural atmospheric CO_2 reservoir has $\delta^{13}\text{C} \approx -7\text{‰}$ when in isotopic equilibrium with marine HCO_3^- and CaCO_3 (Ohmoto, 1986). CO_2 from burning of fossil-fuel and biogenic materials has $\delta^{13}\text{C} \approx -26\text{‰}$ (Hoefs, 1980). Mixing these two CO_2 components with the ratio 21% CO_2 from fossil fuel burning + 79% "natural" CO_2 should give a $\delta^{13}\text{C}$ of the present atmospheric CO_2 of approximately -11‰ .

* Footnote; see last page of the manuscript.

Keeling et al. (1989) have reported $\delta^{13}\text{C}$ of atmospheric CO_2 over the last decades. $\delta^{13}\text{C}$ reported for atmospheric CO_2 was -7.489‰ in December 1978, decreasing to -7.807‰ in December 1988, values close to that of the natural atmospheric CO_2 reservoir, far from the $\delta^{13}\text{C}$ value of -11‰ expected from the IPCC model. Hence the IPCC model is not supported by $^{13}\text{C}/^{12}\text{C}$ evidence.

Segalstad (1992, 1993) has by isotope mass balance considerations calculated the atmospheric CO_2 lifetime and the amount of fossil fuel CO_2 in the atmosphere. The December 1988 atmospheric CO_2 composition was computed for its 748 GT C total mass and $\delta^{13}\text{C} = -7.807\text{‰}$ for 3 components: (1) natural fraction remaining from the pre-industrial atmosphere; (2) cumulative fraction remaining from all annual fossil-fuel CO_2 emissions (from production data); (3) carbon isotope mass-balanced natural fraction. The masses of the components were computed for different atmospheric lifetimes of CO_2 .

The calculations show how the IPCC's (Houghton et al., 1990) atmospheric CO_2 lifetime of 50-200 years only accounts for half the mass of atmospheric CO_2 . However, the unique result fits an atmospheric CO_2 lifetime of ~ 5 (5.4) years, in agreement with numerous ^{14}C studies compiled by Sundquist (1985) and chemical kinetics (Stumm & Morgan, 1970). The mass of all past fossil-fuel and biogenic emissions remaining in the current atmosphere was in December 1988 calculated to be ~ 30 GT C or less, i.e. maximum $\sim 4\%$, corresponding to an atmospheric CO_2 concentration of ~ 14 ppmv.

This small amount of anthropogenic atmospheric CO_2 probably contributes less than half a Watt/m^2 of the $146 \text{ W}/\text{m}^2$ "Greenhouse Effect" of a cloudless atmosphere, contributing to less than half a degree C of radiative heating of the lower atmosphere.

The implication of the ~ 5 year lifetime is that ~ 135 GT C ($\sim 18\%$) of the atmospheric CO_2 pool is exchanged each year. This is far more than the ~ 6 GT C in fossil fuel CO_2 now contributed annually to the atmosphere.

The isotopic mass balance calculations show that at least 96% of the current atmospheric CO_2 is isotopically indistinguishable from non-fossil-fuel sources, i.e. natural marine and juvenile sources from the Earth's interior. Hence, for the atmospheric CO_2 budget, marine equilibration and degassing, and juvenile degassing from e.g. volcanic sources, must be much more important, and burning of fossil-fuel and biogenic materials much less important, than assumed by the authors of the IPCC model (Houghton et al., 1990).

6. Conclusions

Water vapor is the most important "greenhouse gas". Man's contribution to atmospheric CO_2 from the burning of fossil fuels is small, maximum 4% found by carbon isotope mass balance calculations. The "Greenhouse Effect" of this contribution is small and well within natural climatic variability. The amount of fossil fuel carbon is minute compared to the total amount of carbon in the atmosphere, hydrosphere, and lithosphere. The atmospheric CO_2 lifetime is about 5 years. The ocean will be able to absorb the larger part of the CO_2 that Man can produce through burning of fossil fuels. The IPCC CO_2 global warming model is not supported by the scientific data. Based on geochemical knowledge there should be no reason to fear a climatic catastrophe

because of Man's release of the life-governing CO₂ gas. The global climate is primarily governed by the enormous heat energy stored in the oceans and the latent heat of melting of the ice caps, not by the small amount of heat that can be absorbed in atmospheric CO₂; hence legislation of "CO₂ taxes" to be paid by the public cannot influence on the sea level and the global climate.

Acknowledgements: Technological Oriented Studies, University of Oslo, for financial support; Dr. Zbigniew Jaworowski for scientific discussions and contributions.

References

- Bolin, B. & Keeling, C.D. (1963): Large-scale atmospheric mixing as deduced from the seasonal and meridional variations of carbon dioxide. *Journal of Geophysical Research* 68 (13), 3899-3927.
- Butler, J.N. (1982): *Carbon dioxide equilibria and their applications*. Addison-Wesley Publishing Company, 259 pp.
- Druffel, E.R.M. & Williams, P.M. (1990): Identification of a deep marine source of particulate organic carbon using bomb ¹⁴C. *Nature* 347, 172-174.
- Drummond, S.E., Jr. (1981): *Boiling and mixing of hydrothermal fluids: chemical effects on mineral precipitation*. Ph.D. thesis, The Pennsylvania State University, 381 pp.
- Faure, H. (1990): Changes in the global continental reservoir of carbon. *Paleontology, Paleogeography, Paleoecology (Global Planetary Change Section)* 82, 47-52.
- Hoefs, J. (1980): *Stable isotope geochemistry, 2nd ed. Minerals and Rocks*, 9. Springer-Verlag, 208 pp.
- Heyke, H.-E. (1992 a): Zu den CO₂-Klimakurven aus Eisbohrkernen. *Erdöl und Kohle, Erdgas, Petrochemie vereinigt mit Brennstoff-Chemie* 45 (5), 208-214.
- Heyke, H.-E. (1992 b): Zu den CO₂-Klimakurven aus Eisbohrkernen - Diskussion. *Erdöl und Kohle, Erdgas, Petrochemie vereinigt mit Brennstoff-Chemie* 45 (9), 360-362.
- Heyke, H.-E. (1992 c): Gasblasen im Eis sind brüchiges Fundament für die CO₂-Steuer. *Fusion* 13 (3), 32-39.
- Holland, H.D. (1984): *The chemical evolution of the atmosphere and oceans*. Princeton University Press, 582 pp.
- Houghton, J.T., Jenkins, G.J. & Ephraums, J.J. (Eds.) (1990): *Climate Change. The IPCC Scientific Assessment. Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, 365 pp.
- Jaworowski, Z., Segalstad, T.V. & Hisdal, V. (1992 a): Atmospheric CO₂ and global warming: a critical review; 2nd revised edition. *Norsk Polarinstitutt, Meddelelser [Norwegian Polar Institute, Memoirs]* 119, 76 pp.

- Jaworowski, Z., Segalstad, T.V. & Ono, N. (1992 b): Do glaciers tell a true atmospheric CO₂ story? *Science of the Total Environment* 114, 227-284.
- Keeling, C.D., Bacastow, R.B., Bainbridge, A.E., Ekdahl, C.A. Jr., Guenther, P.R., Waterman, L.S. & Chin, J.F.S. (1976): Atmospheric carbon dioxide variations at Mauna Loa Observatory, Hawaii. *Tellus* 28 (6), 538-551.
- Keeling, C.D., Bacastow, R.B., Carter, S.C., Whorf, T.P., Heimann, M., Mook, W.G. & Roeloffzen, H. (1989): A three-dimensional model of atmospheric CO₂ transport based on observed winds: 1. Analysis of observational data. In Peterson, D.H. (Ed.): Aspects of climate variability in the Pacific and the Western Americas. *American Geophysical Union, Geophysical Monograph* 55, 165-236.
- Kerr, R.A. (1992): Fugitive carbon dioxide: it's not hiding in the ocean. *Science* 256, 35.
- Kuo, C., Lindberg, C. & Thomson, D.J. (1990): Coherence established between atmospheric carbon dioxide and global temperature. *Nature* 343, 709-714.
- MacIntyre, R. (1970): Why the sea is salt. *Scientific American* 223 (5), 104-115.
- Krauskopf, K.B. (1979): *Introduction to geochemistry; 2nd edition*. McGraw-Hill, Inc., 617 pp.
- Maier-Reiner, E. & Hasselmann, K. (1987): Transport and storage of CO₂ in the ocean - an inorganic ocean-circulation carbon cycle model. *Climate Dynamics* 2, 63-90.
- Oerlemans, J. & van der Veen, C.J. (1984): *Ice sheets and climate*. D. Reidel Publishing Company, 217 pp.
- Ohmoto, H. (1986): Stable isotope geochemistry of ore deposits. *Reviews in Mineralogy* 16, 491-559.
- Peixoto, J.P. & Oort, A.H. (1992): *Physics of climate*. American Institute of Physics, 520 pp.
- Ramanathan, V., Barkstrom, B.R. & Harrison, E.F. (1989): Climate and the Earth's radiation budget. *Physics Today* 42 (5), 22-32.
- Raval, A. & Ramanathan, V. (1989): Observational determination of the greenhouse effect. *Nature* 342, 758-761.
- Revelle, R. & Suess, H.E. (1957): Carbon dioxide exchange between atmosphere and ocean and the question of an increase of atmospheric CO₂ during the past decades. *Tellus* 9, 18-27.
- Segalstad, T.V. (1990): Temperatur og CO₂. *Teknisk Ukeblad* 137 (17; yellow part), 4-5.
- Segalstad, T.V. (1992): The amount of non-fossil-fuel CO₂ in the atmosphere. *American Geophysical Union, Chapman Conference on Climate, Volcanism, and Global Change, March 23-27, 1992, Hilo, Hawaii. Abstracts*, 25.

Segalstad, T.V. (1993): Stable isotope geochemistry applied to paleoclimatological and greenhouse gas problems. *1st International Symposium on Applied Isotope Geochemistry (AIG-1)*, Aug. 29 - Sept. 3, 1993, Geiranger, Norway. *Program and Abstracts, Institute for Energy Research IFE/KR/E-93/007*, 95-96.

Segalstad, T.V. & Jaworowski, Z. (1991): CO₂ og globalt klima. *Kjemi 51 (10)*, 13-15.

Skirrow, G. (1975): The dissolved gases - carbon dioxide. In Riley, J.P. & Skirrow, G. (Eds.): *Chemical oceanography, Vol. 2; 2nd edition*. Academic Press, 1-192.

Stumm, W. & Morgan, J.J. (1970): *Aquatic chemistry. An introduction emphasizing chemical equilibria in natural waters*. John Wiley & Sons, 583 pp.

Sundquist, E.T. (1985): Geological perspectives on carbon dioxide and the carbon cycle. In Sundquist, E.T. & Broecker, W.S. (Eds.): *The carbon cycle and atmospheric CO₂: natural variations Archean to present. American Geophysical Union, Geophysical Monograph 32*, 5-59.

Takahashi, T. (1961): Carbon dioxide in the atmosphere and in Atlantic ocean water. *Journal of Geophysical Research 66 (2)*, 477-494.

Volk, T. & Liu, Z. (1988): Controls of CO₂ sources and sinks in the Earth scale surface ocean: temperature and nutrients. *Global Biogeochemical Cycles 2*, 73-89.

Footnote to the manuscript:

*

$$\delta^{13}\text{C} = \frac{(^{13}\text{C}/^{12}\text{C})_{\text{sample}} - (^{13}\text{C}/^{12}\text{C})_{\text{standard}}}{(^{13}\text{C}/^{12}\text{C})_{\text{standard}}} \cdot 1000 \text{ ‰}$$

where the reference standard used here is PDB (Pee Dee Belemnite) CaCO_3 .

Biography of Tom Victor Segalstad

Born in Norway in 1949. University degrees (natural sciences with geology) from The University of Oslo. Has conducted university research, publishing, and teaching in geochemistry, mineralogy, petrology, volcanology, structural geology, ore geology, and geophysics at The University of Oslo, Norway, and The Pennsylvania State University, USA. At present keeping professional positions as Associate Professor of Geochemistry at The University of Oslo; Head of The Mineralogical-Geological Museum at The University of Oslo; and Director of The Natural History Museums and Botanical Garden of The University of Oslo. Member of different international and national professional working groups and committees.

Oslo, December 1995.

Manuscript for the chapter printed on pages 41-50 in the book:

Emsley, J. (Ed.) 1996: *The Global Warming Debate. The Report of the European Science and Environment Forum*. Bourne Press Ltd., Bournemouth, Dorset, UK. (ISBN 0952773406). 288 pp.

